

PATENT SPECIFICATION

843,135

NO DRAWINGS.

Inventors:—SIEGFRIED GOTTFRIED and LILY BAXENDALE.



Date of filing Complete Specification : June 6, 1958.

Application Date : July 16, 1957. No. 26453/59.
(Divided out of No. 843,133).

Complete Specification Published : Aug. 4, 1960.

Index at Acceptance :—Classes 2(3), B4(A1 : A2 : A4 : M), C2A(1 : 5), C2B3(A1 : E : G1), C(2R15 : 3A16) ; and 81(1), B1(C : F : H : K : Q : R1 : R2 : S : Z), B2(C : F : H : K : Q : R1 : R : 2 S : Z).

International Classification :—B61k. C07d, g.

COMPLETE SPECIFICATION.

Glycyrrhetic Acid Salts.

5 We, BIOREX LABORATORIES LIMITED, a British Company, organised under the laws of Great Britain, of 47/51 Exmouth Street, Rosebery Avenue, London, E.C.1, do hereby declare the invention, for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—

10 This invention comprises improvements in and relating to pharmacological compounds, and more particularly new derivatives of glycyrrhetic acid.

15 Glycyrrhetic acid is obtainable from liquorice root. In Specifications Nos. 798,655 and 799,415 we have described and claimed compositions containing glycyrrhetic acid and having a pronounced effect in suppressing inflammation. The said compositions are suitable for topical application. Glycyrrhetic acid being only sparingly soluble in water.

20 We have now found that the organic salts of glycyrrhetic acid also have a good effect in suppressing inflammation, and yet are more soluble generally, especially in body fluids than glycyrrhetic acid, and are, therefore, more suitable for systemic use than is glycyrrhetic acid. It is to be understood that the new compounds are prepared from non-toxic bases which are compatible with mammals.

25 The new organic salts of glycyrrhetic acid are prepared by methods in use or described in the literature for the conversion of organic acids to their salts with organic bases. Thus, they may be prepared by the interaction of the organic base with glycyrrhetic acid, preferably in aqueous or organic media; in some cases an elevated temperature is useful in order to accelerate the salt formation.

40

45 These new compounds have a good effect in suppressing inflammation and may be used in compositions with known additives, such as with inert carriers, to form, for example, an ointment, powder, or emulsion, and may also be compounded with anti-causative agents, as is the case with glycyrrhetic acid itself (e.g. as described in Specifications Nos. 798,655, 799,415 and 26332/57 (Serial No. 843,134)). Further the new compounds may be dissolved in suitable solvents, such as water, normal saline or oils, and so make compositions suitable for, for example oral, sub-cutaneous, intramammary, intra-muscular, intra-articular, intra-peritoneal and intravenous use.

50
55
60 Medical, pharmacological, and veterinary tests and trials have been carried out with these new compounds on human beings, small and large animals, as well as pharmacological trials using rats, mice, guinea pigs, rabbits and cats. The new compounds have shown that the hereinafter-described derivatives are active in suppressing inflammation, for example pharmacological tests carried out as is more particularly described below:

65
70 These new compounds heal artificial lesions produced on the skin of rabbits, whether from an external cause, or from intradermal injections of irritant substances. When applied locally, these new compounds cause rapid subsidence of any inflammation produced by the introduction of irritant substances into the eye of the rabbit.

When applied by systemic injection or by oral administration these new compounds depress the formation of granuloma tissue induced by subcutaneously-implanted cotton wool pellets in rats in the test described by Meier, R., Schuler, W., and Desaulles, P., *Experientia*, 1950, 6, 469. These new derivatives depress the formation of inflammatory exudate and of the granulomatous membrane in the granuloma pouch test described by Selye, H., *Britt. med. J.*, 1949, 2, 1129.

When injected systemically into B.C.G.-infected guinea pigs, the new derivatives suppress the reaction to intra-dermally-injected tuberculin in the test described by Long, D. A., and Miles, A. A., *Lancet*, 1950, 1, 492.

In addition, when injected parenterally or administered orally, the new derivatives have a mild depressant action in mice and potentiate the actions of central nervous system depressant drugs such as hexobarbitone.

The new derivatives have mild analgesic and antipyretic actions.

These new compounds are of value in combating inflammatory conditions of all types of denominations, such as inflammatory conditions of the skin, eye, ear, nose, mouth, dental cavities, genitals, rheumatic conditions, rectal conditions, inflammatory and ulcerative conditions of the digestive system, ulcerative colitis, allergic conditions, vaginitis, vulvitis, dysmenorrhoea, metritis, leukorrhoea, mastitis, and other inflammatory processes whether they be primary or of secondary cause, or the result of such cause.

These new compounds have also been found valuable in the treatment of severe emergencies, in which a "shock-like" state occurs, by virtue of their high solubility in water.

These new compounds have also been found of value when injected into inflamed joints and it was found that symptomatic relief was obtained.

In addition, these new compounds are of value in a number of illnesses where a mild sedative, analgesic, or antipyretic may be indicated.

The glycyrrhetic acid derivatives according to the present invention inhibit steroids and their metabolism, such as the usual hormonal secretions of glands, by reduction of glandular activity. They are useful in the treatment of anogenital diseases, such as vulvitis, vaginitis, ulcerations of the vaginal basin and cervix of the uterus, either alone or in association with leucorrhoea, menstrual disfunctions, ano-genital pruritis, and in the treatment of diseases, such as carcinomas, in which the inhibition of steroids or the reduction of hormonal glandular secretion is of therapeutic value, as well as inflammatory

and ulcerative conditions of the digestive system.

The new compounds have a potentiating effect demonstrating synergism with antibiotic drugs, (such as Neomycin), keratoplastic drugs, (such as Coal Tar), keratolytic drugs, (such as Salicylic Acid), analgesics, antiseptics, bacteriocides, chemotherapeutics, bacteriostatics, anti-histaminics, sedatives, fungicides, insecticides, corticosteroids and xanthoglabrol.

Synergism has also been demonstrated in conjunction with corticosteroids, such others as hydrocortisone, prednisone and prednisolone, in the replacement therapy of adrenalectomised patients, as well as in Addison's disease, disseminated lupus erythematosus and acute bronchial asthma.

As has already been indicated, these new derivatives can be incorporated in various therapeutic forms, such as, ointments, solutions, injections, emulsions, suspensions, pastes, cones, cerates, paints, powders, and implants, all in conjunction with suitable carriers.

The following examples are given for the purpose of illustrating the invention:—

EXAMPLE 1.

Piperazine salt of glycyrrhetic acid. 2.0 grams of piperazine hexahydrate were dissolved in 200 cc. of boiling water. 9.0 grams of glycyrrhetic acid were added, as an aqueous paste, and then 200 cc. of rectified spirit. On boiling on a water bath, practically all the acid dissolved, and the suspension was filtered and the filtrate evaporated on the boiling water bath. Removal of water was completed by vacuum-drying over solid potassium hydroxide, and the solid piperazine salt of glycyrrhetic acid obtained was then readily ground to a fine powder. This salt is more soluble in water than is glycyrrhetic acid.

EXAMPLE 2.

N-methyl glucamine salt of glycyrrhetic acid.

A solution of 8 grams of N-methyl-glucamine in 50 cc. of water was added to 19 grams of glycyrrhetic acid, previously ground to a paste with 100 cc. of water. After warming to about 50° C. and shaking for a few minutes, the suspension was filtered and the filtrate evaporated on a water bath. The residue, which was the N-methyl-glucamine salt of glycyrrhetic acid, was further freed from water by drying in a vacuum over concentrated sulphuric acid. The salt is very readily soluble in water, and is a white powder.

The triethanolamine salt and salts of other organic bases may be prepared in a similar manner.

EXAMPLE 3.

An ointment was prepared by dissolving 2% by weight of the piperazine salt of glycyrrhetic acid and by 5% by weight of neomycin sulphate in a "Vaseline" (Registered Trade Mark) base. The base can also be made from 8% by weight of non-ionic emulsifying wax B.P.C. and 16% by weight of liquid paraffin in water.

0.1 g. "Edipas" (Registered Trade Mark), which is an edible cellulose derivative, in normal saline to make 20 ml.

Further ointments, emulsions, lotions, suppositories, tablets, injections and pessaries were prepared in a manner similar to those given in Examples 3—8 in which the glycyrrhetic acid derivative was the sole active ingredient present.

EXAMPLE 4.

Ointments were prepared as described in Example 3 in which the neomycin sulphate was replaced by an equal amount of centrimide.

EXAMPLE 5.

An emulsion was prepared by dissolving 0.5% by weight of the N-methylglucamine salt of glycyrrhetic acid, 0.5% by weight of hexachlorophene, 5% by weight of "Lanbritol" (Registered Trade Mark), 10% by weight of glycerol and 20% by weight of polyethylene glycol in water.

EXAMPLE 6.

A lotion was prepared by dissolving 1% by weight of the piperazine salt of glycyrrhetic acid, 0.5 by weight of hydrocortisone alcohol, 0.5% by weight of coal tar fractions and 10% by weight of polyethylene glycol (molecular weight 600) in water. This example illustrates the replacement of hydrocortisone in a known preparation by a glycyrrhetic acid derivative to give a preparation with an enhanced anti-inflammatory effect.

EXAMPLE 7.

A suppository was prepared by dispersing 1% by weight of the triethanolamine salt of glycyrrhetic acid, 2% by weight of benzocaine and 5% by weight of 2:4- or 4:4- (diacetoxydiphenylmethyl) pyridine in a suppository base (British Pharmaceutical Codex).

EXAMPLE 8.

An injection was prepared containing 0.25 g. of the N-methylglucamine salt of glycyrrhetic acid 0.048 g. methyl *p*-hydroxybenzoate, 0.08 g. "Tween" (Registered Trade Mark), which is a wetting agent and

WHAT WE CLAIM IS:—

1. Salts of glycyrrhetic acid with organic bases.
2. Piperazine salt of glycyrrhetic acid.
3. N-methylglucamine salt of glycyrrhetic acid.
4. Triethanolamine salt of glycyrrhetic acid.
5. Process for the production of organic salts of glycyrrhetic acid, wherein the derivatives are prepared from glycyrrhetic acid by methods known *per se* for the conversion of organic acids to their salts with organic bases.
6. Process for the production of organic salts of glycyrrhetic acid, substantially as hereinbefore described and with reference to either of Examples 1 and 2.
7. Organic salts of glycyrrhetic acid, whenever prepared by the process according to Claim 5 or 6.
8. Pharmaceutical and therapeutic compositions containing as active ingredient one or more glycyrrhetic acid derivatives according to any of Claims 1—4 and 7, alone or in admixture with one or more antibiotics, keratoplastics, keratolytics, analgesics, bacteriocides, chemotherapeutics, bacteriostatics, corticosteroids, xanthoglabrol, anti-histaminics, sedatives, fungicides and/or insecticides, and an inert pharmaceutical carrier.
9. Pharmaceutical and therapeutic compositions substantially as hereinbefore described with reference to any of Examples 3—8.

CARPMAELS & RANSFORD,
Agents for Applicants,
24 Southampton Buildings,
Chancery Lane, London, W.C.2.

PROVISIONAL SPECIFICATION.

Glycyrrhetic Acid Salts.

We, BIOREX LABORATORIES LIMITED, a British Company, organised under the laws of Great Britain, of 47/51 Exmouth Street, Rosebery Avenue, London, E.C.1, do hereby declare this invention to be described in the following statement:—

This invention comprises improvements in and relating to pharmacological com-

pounds, and more particularly new derivatives of glycyrrhetic acid.

Glycyrrhetic acid is obtainable from liquorice root. In Specification Nos. 8183/56 and 8184/56 (Serial Nos. 798,655 and 799,415) we have described and claimed compositions containing glycyrrhetic acid and having a pronounced effect in suppress-

ing inflammation. The said compositions are suitable for topical application, glycyrrhetic acid being only sparingly soluble in water.

5 We have now found certain derivatives of glycyrrhetic acid, which also have a good effect in suppressing inflammation, and yet are more soluble generally, especially in body fluids, than glycyrrhetic acid, and are therefore, more suitable for
10 systematic use than in glycyrrhetic acid.

According to the invention, there are provided, as new compounds, organic salts of glycyrrhetic acid. These new compounds have a good effect in suppressing inflammation and may be used in compositions with known additives, e.g. with inert carriers to form, e.g. an ointment, powder, or emulsion and also may be compounded with anti-causative agents, as is the case with glycyrrhetic acid itself, e.g. as described in Specification Nos. 8183/56 and 8184/56 (Serial Nos. 798,665 and 799,415). Further the new compounds may be dissolved in
25 suitable solvents, oral, e.g. water, normal saline, etc., and so make compositions suitable for e.g., sub-cutaneous, intra-muscular, intra-articular, intra-peritoneal and intra-venous use.

30 The following examples are given for the purpose of illustrating the invention, showing examples of new salts with organic bases.

For instance, new salts of glycyrrhetic acid not described hitherto have been prepared.

EXAMPLE 1.

Piperazine salt of glycyrrhetic acid.

2.0 grams of piperazine hexahydrate were dissolved in 200 cc. of boiling water. 9.0
40 grams of glycyrrhetic acid were added, as an aqueous paste, and then 200 cc. of rectified spirit. On boiling on a water bath, practically all the acid dissolved, and the
45 suspension was filtered and the filtrate evaporated on the boiling water bath. Removal of water was completed by vacuum-drying over solid potassium hydroxide, and the solid piperazine salt of glycyrrhetic acid obtained was then readily ground to
50 a fine powder. This salt is more soluble in water than is glycyrrhetic acid.

EXAMPLE 2.

55 N-methyl glucamine salt of glycyrrhetic acid.

A solution of 8 grams of methyl glucamine in 50 cc. of water was added to 19 grams of glycyrrhetic acid, previously ground to a paste with 100 cc. of water. After warming to about 50° C. and shaking for a few
60 minutes, the suspension was filtered and the filtrate evaporated on a water bath. The residue, which was the N-methyl glucamine salt of glycyrrhetic acid, was further freed

65 from water by drying in a vacuum over concentrated sulphuric acid. The salt is very readily soluble in water, and is a white powder.

The triethanolamine salt and salts of other organic bases may be prepared in a
70 similar manner.

Medical, pharmacological, and veterinary tests and trials have been carried out with these new compounds on human beings, small and large animals, as well as
75 pharmacological trials using rats, mice, guinea pigs, rabbits and cats. The new compounds have shown that the afore-described derivatives are active in suppressing inflammation, for example pharmacological tests carried out as is more
80 particularly described below:—

These new compounds heal artificial lesions produced on the skin of rabbits, whether from an external cause, or from
85 intradermal injections of irritant substances.

When applied locally, these new compounds cause rapid subsidence of any inflammation produced by the introduction of irritant substances into the eye of the rabbit.
90

When applied by systemic injection or by oral administration these new compounds depress the formation of granuloma tissue induced by sub-cutaneously implanted cotton wool pellets in rats in the test described by
95 Meier, R., Schuler, W., and Desaulles, P., *Experientia*, 1950, 6, 469.

These new derivatives depress the formation of inflammatory exudate and of the granulomatous membranes in the granuloma
100 pouch test described by Selye, H., *Brit. med. J.*, 1949, 2, 1129.

When injected systemically into B.C.G.-infected guinea pigs, the new derivatives suppress the reaction to intradermally-
105 injected tuberculin in the test described by Long, D. A., and Miles, A. A., *Lancet*, 1950, 1, 492.

In addition, when injected parenterally or administered orally, the new derivatives
110 have a mild depressant action in mice and potentiate the actions of central nervous system depressant drugs such as hexobarbitone.

The new derivatives have mild analgesic
115 and antipyretic actions.

These new compounds are of value in combating inflammatory conditions of all types and denominations, such as inflammatory conditions of the skin, eye, ear,
120 nose, mouth, dental cavities, genitals, rheumatic conditions, rectal conditions, inflammatory and ulcerative conditions of the digestive system, ulcerative colitis, allergic conditions, vaginitis, vulvitis, dysmenorrhoea, metritis, leukorrhoea, mastitis, and other
125 inflammatory processes whether they be primary or of secondary cause, or the result of such cause.

- These new compounds have also been found valuable in the treatment of severe emergencies, in which a "shock-like" state occurs, by virtue of their high solubility in water. 5
- These new compounds have also been found of value when injected into inflamed joints and it was found that symptomatic relief was obtained.
- 10 In addition, these new compounds are of value in a number of illnesses where a mild sedative, analgesic, or anti-pyretic may be indicated.
- 15 The new compounds have a potentiating effect demonstrating synergism with anti-biotic drugs, (such as Neomycin), keratoplastic drugs, (such as Coal Tar), keratolytic drugs, (such as Salicylic Acid), antiseptics, bacteriocides, chemotherapeutics, bacteriostatics, anti-histaminics, sedatives, fungicidal, and insecticidal. 20
- Synergism has also been demonstrated in conjunction with corticosteroids, such others as hydrocortisone, prednisone, prednisolone, etc., in the replacement therapy of adrenalectomised patients, as well in Addison's disease, disseminated lupus erythematosus, acute bronchial asthma, etc. 25
- As has already been indicated, these new derivatives can be incorporated in various therapeutic forms, such as, ointments, solutions, injections, emulsions, suspensions, pastes, cones, cerates, paints, powders, all in conjunction with suitable carriers. 30
- CARPMAELS & RANSFORD,
Agents for Applicants,
24 Southampton Buildings,
Chancery Lane, London, W.C.2.

Abingdon : Printed for Her Majesty's Stationery Office, by Burgess & Son (Abingdon), Ltd.—1960.
Published at The Patent Office, 25, Southampton Buildings, London, W.C.2,
from which copies may be obtained.

THIS PAGE BLANK (USPTO)